Proposals to Revise Soil Taxonomy

Craig Ditzler

National Leader, Soil Classification and Standards

Micheal L. Golden

MLRA Leader Region 9/Texas State Soil Scientist

NCSS South Region Soil Taxonomy Committee

MLRA

Office	Co-Chair's	<u>Start</u>	End
• MO-9	Mike Golden	June 2000	June 2002
• MO-1	4 Roy Vick	June 2002	June 2004
• MO-1	5 Charles Love	June 2004	June 2006
• MO-1	6 Charles Fultz	June 2006	June 2008
• MO-1	8 Bill Craddock	June 2008	June 2010
• NSSC	Craig Ditzler	Permanent	Co-Chair 1/

1/ (National Leader - Classification & Standards)

NCSS South Region Soil Taxonomy Committee

			<u>Start</u>	End
•	Mike Verpraska	s - Member	Jan 1999	Dec 2001
•	Larry West	- Member	Jan 2000	Dec 2002
•	Moye Rutledge	- Member	Jan 2001	Dec 2003
•	Joey Shaw	- Member	Jan 2002	Dec 2004
•	Mike Golden	- Co-Chair	June 2000	June 2002
•	Roy Vick	- Co-Chair	June 2002	June 2004
•	Craig Ditzler	- Co-Chair	Permanent	

18 Proposals to Amend Soil Taxonomy

- Letter from Craig Ditzler Dated 1/23/2002
 - Each NCSS Region is to have comments and recommendations by August 30, 2002
 - Respond to each proposal with following:
 - Accept as is
 - Accept with revision (explain revisions)
 - Return to originator for further consideration
 - Reject (explain)

South Region Soil Taxonomy Committee

Region	Proposal
Northeast	1
South	2
Midwest	3-7
West	8-12
NHQ & NSSC Staff	13-18

1) Subaqueous Subroups George Demas, MD (Deceased)

- Aquents that are "Permanantly submersed"
- Key out first in each great group
- Soil = "water not too deep for rooted plants"
- Proposed for use in:
 - Endoaquents, Fluvaquents, Hydraquents
 - Psammaquents, Sulfaquents

1) Subaqueous Subroups

George Demas, MD (Deceased)

13 Subgroups are proposed

- All in the Aquents
- Example:
 - "Endoaquents which have sulfidic materials within 100 cm of the mineral soil surface and are permanently submersed"
 - » Subaquic Sulfic Endoaquents

Footnote: Some OSD's will be affected in South Region but extent unknown

2) Humic subgroups in some Inceptisols Roy Vick, Bill Craddock, Gene Mayhugh

- Many former Umbrepts and Ochrepts went to Udepts
- Ochrepts with dark (but thin) surfaces wound up in Typic subgroups
- Rename present Humic subgroups to Umbric
- Establish (3) new subgroups for thin, dark ochrics Humic Dystrudepts, Fluventic Humic Dystrudepts

 Humic Fragiudepts

Footnote: Unknown number of OSD's affected in South Region Roy will cover this proposal in more detail

3) Fragipan Definition Dr. Don Franzmeier

- Dense calcareous tills differ from fragipans in genesis, kinds and rates of weathering, plant nutrient chemistry, and reactions with leaching chemicals.
- Remove midwestern dense tills from fragipans by: Adding a 6th required characteristic to the Fragipan: "Is not effervescent"
- Presumably would be densic material.

Footnote: Unknown number of OSD's affected in South Region

4) Miscellaneous Changes Bismark MO

- Natrustalfs Add: Aridic Glossic subgroup
- Argiudolls, Hapludolls, and Paleudolls
 - Add: Aquic Pachic subgroup & change keying order
- Hapludolls Add: Pachic Vertic subgroup

Footnote: OSD's affected in South Region

- No changes in Natrustalfs
- Aquic Pachic Argiustolls (LA- Armistead; TX- Benklin, Faddin, & Waskom; OK- Garton)
- Aquic Pachic Paleustolls (OK Burwell & Okemah)
- Pachic Vertic Hapludolls (AR- Desha)

Miscellaneous Changes Bismark MO

- Argiustolls -Add subgroups
 - Aquertic
 - Pachic Vertic
 - Pachic Udertic

Footnote: OSD's affected in South Region

- Aquertic (OK- Foraker, Tabler, & Westum; TX- Flatonia)
- Pachic Udertic (OK- Brewer and Brewless; TX- Elmenwood)
- Pachic Vertic (TX- Coy, Culp, Elmendorf, Laparita, Loftan & Popia)

Miscellaneous Changes Bismark MO

- Haplustolls Add subgroups:
 - Aridic Lithic
 - Aquertic
 - Pachic Udertic
 - Pachic Vertic

Footnote: OSD's affected in South Region

- Aridic Lithic Several unspecified OSD's affected
- Aquertic No OSD's affected
- Pachic Udertic (TX- Divot, Eloso, Nukrum, & Rioconcho)
- Pachic Vertic No OSD's affected

Miscellaneous Changes Bismark MO

- Argiustolls, Calciustolls, Haplustolls
 - Reorder keys
 - From: Pachic Aquic Oxyaquic
 - To: Aquic Oxyaquic Pachic
- Paleustolls
 - Reorder keys
 - From: Pachic Aquic
 - To: Aquic Pachic

Footnote: No changes to OSD's recognized in South Region

4(b) - Pachic Subgroup Definitions Gene Mayhugh

• Clarify definition to indicate that the requirement for texture finer than LFS applies to the mollic epipedon

4 (c, d) Miscellaneous Bismarck MO

- Dystrustepts
 - Add: Torrertic, Vertic, & Aridic subgroups
- Natrargids
 - Add: Ustertic, & Xerertic subgroup

Footnote: No changes to OSD's in South Region

5) Aquic: Hapludolls, Argiudolls, Paleudolls C.J. Heidt/G. Mayhugh

- Carbon content of sandy soils decreases before color does. (actual mollic thinner than it looks).
 - 1) Allow (aquic) colors below the dark horizon, not immediately below "mollic".

OR

2) Change reference point from base of mollic to

"< 100 cm of mineral surface" or

"≤ 75 cm of mineral surface"

Footnote:

- Argiudolls (TX- Chireno & Labelle; OK- Summit)
- Paleudolls (OK- Burwell, Choteau, & Okema)
- Haplustolls No changes to OSD's

6) Miscellaneous Changes Indianapolis MO

• A) Add: Mollic Oxyaquic Hapludalfs

- B) Spodic subgroups:
 - Allow: Albic underlain by horizon 1 hue darker

Footnote:

- A) Mollic Oxyaquic Hapludalf (LA- Hendrick) only OSD affected
- B) No OSD's affected in South Region

6) Miscellaneous Changes Indianapolis MO

• C) Hapludolls

Rearrange Keying Order:

 Move Fluventic to just below Fluvaquentic and ahead of Pachic and Oxyaquic suborders

Footnote: No OSD's affected in South Region

6) Miscellaneous Changes Indianapolis MO

- D) Endoaquepts
 - Add: Fluventic Subgroup.
- E) Haplorthods
 - Add: Lamellic Oxyaquic Subgroup

Footnote:

- D) Fluventic Endoaquepts (MS- Arklabutla; TX- Manco & Pophers)
- E) No OSD's affected in South Region

7) Miscellaneous Changes Indianapolis MO

- Haplorthods
 - Add: Alfic Lamellic subgroup
- Glossudalfs
 - Add: Aquic Arenic & Arenic Oxyaquic subgroups
- Hapludalfs
 - Add: Arenic Oxyaquic subgroup

Footnote: No OSD changes in South Region for Haplorthods or Glossudalfs

• Arenic Oxyaquic Hapludalfs - (AR- Bulltown; FL- Chiefland)

8) Depth of Eutrocryepts Lakewood MO

• Revise keys to require $\geq 60\%$ BS between 25 and 75 cm from soil surface (current criteria)

But add:

"or immediately above a root limiting layer if at a shallower depth."

Footnote: No OSD's affected in South Region

9) Torripsamments Reno MO

Add: Oxyaquic subgroup

 Have water within 150 cm for 20 consecutive or 30 cumulative days

Footnote: No OSD's affected in South Region

10) Miscellaneous Changes Portland MO

 Several changes to Cryic and Andic soils in far west

Footnote: No changes to OSD's in South Region

11) Revise Dystric subgroups Reno MO

- Several changes to
 - Xerorthents and Xeropsamments in western US

Footnote: No changes to OSD's in South Region

12) Strongly Contrasting Particle-size Classes Portland MO

- Add 2 new classes:
 - Loamy over ashy or ashy-pumiceous
 - Medial-skeletal over loamy-skeletal
- Added Definition of Diatomaceous mineralogy class

Footnote: No known OSD's affected in South Region

13) New Greatgroup and Subgroups for Aquerts Hari Eswarin - NHQ

- Add Greatgroup: Sulfaquerts.
 - Salic subgroup
 - Sulfic subgroup
 - Typic subgroup

Footnote: Only for soils in Thailand

14) Add Spodic Subgroups NSSC - Hipple/Engel

- Spodic Haplocryands
- Spodic Vitricryands

Footnote: No OSD's in South Region are affected

15) Clarify definitions of "Resistant" and "Weatherable" minerals. NSSC - Engel

- Resistant: Add "iron oxides, kaolinite, gibbsite, chalcedony... as well as others" to list in text.
- Weatherable:
 - For clay minerals, add allophane to list of "also included".
 - For silt/sand size: Add dolomite to list. Clarify that calcite, carbonates, gypsum, and halite are NOT included, due to mobility and recharge.

Footnote: Several OSD's in SW part of South Region could be affected

16) Key to Mineralogy NSSC - Lynn

- Several items, specify percent by weight.
- Several items, specify iron is Fe₂O₃.
- Kaolinitic rewritten (no reference to nonexpanding 2:1, gibbsite, or halloysite).
- Halloysitic require < 10% smectite.
- Several items, specify extractions from fine-earth fraction.

Key to Mineralogy NSSC - Lynn

- Glassy add "glass-coated grains, glassy materials, and glassy aggregates."
 delete size fraction.
- Gibbsitic delete reference to "hydrated aluminum oxides"
- Parasesquic criteria revised, gibbsite removed.

Key to Mineralogy NSSC - Lynn

- Several items specify thickness criteria applies to mineralogy control section.
- Micaceous & Paramicaceous
 - Reduce required grain count percent.
 - Narrow applicable size range
 from 0.02-2.0 mm to 0.02 0.25 mm.
- Siliceous Allow grain count. Change "extremely durable" to "resistant minerals".

17) Keying Order Gene Mayhugh

Coordinate varying taxa to key in the order:
 Anthraquic - Aquic - Oxyaquic.

Footnote: No OSD's affected in South Region

18) Mollic/Umbric Epipedon Thickness NSSC Engel/Ditzler

- Restore mistakenly omitted criteria regarding carbon with depth.
- Depth requirements rewritten for clarity.

 No substantial change intended.

Footnote: No OSD's affected in South Region

Proposals to Taxonomy

- Give written <u>Comments</u> to any of the South Region NCSS Committee members by August 1, 2002.
- The committee will consolidate comments for proposals and submit them to NSSC for the **South Region**.

Thank You